

TITLE

Are cycle ergometers effective to promote physical activity and physical function among older adults? Short-term effects of a 10-week structured versus need-supportive physical activity program in assisted living facilities.

AUTHORS

Sofie Martien, Filip Boen, Jan Seghers, Christophe Delecluse

INSTITUTIONS

Physical Activity, Sports and Health Research Group, Department of Kinesiology, Faculty of Kinesiology and Rehabilitation Sciences, KU Leuven, Belgium

Acknowledgements

This study was conducted under the authority of the Policy Research Centre on Sports, supported by the Flemish Government.

ABSTRACT

Introduction

The population of adults over the age of 70 is rapidly growing. A large percentage of this population is not regularly involved in physical activity (PA), even though PA can attenuate the impact of aging. The current study examined the short-term effectiveness of two cycle ergometer programs on PA and physical function in older adults (age ≥ 70 yr).

Methods

Flemish older adults living in an assisted living facility were motivated to use a cycle ergometer by assigning them either to (1) a structured PA program (STRUC, $n=35$), which consisted of three times weekly supervised PA sessions, or to (2) a need-supportive PA program (NEED, $n=36$), including four 1-hour individual contact moments. NEED received individually tailored PA coaching based on the Self-Determination Theory. Participants of both STRUC and NEED received an individualized cycle ergometer program. Results were compared with (3) a control group (CON, $n=24$) before (pretest) and immediately after (posttest) the 10-week intervention period with respect to self-reported PA, functional performance (modified Physical Performance Test), exercise capacity (6-minute walk test) and lower body strength (knee-extension test).

Results

The study sample consisted of 22 men and 73 women, with a mean age of 82 ± 6 years. Repeated measurement ANOVA showed a significant time-by-group interaction effect with respect to self-reported moderate-intensity PA (min/week). More specifically, moderate-intensity PA significantly increased in both intervention groups from pre- to posttest (STRUC: +41 min/week, NEED: +44 min/week), whereas no changes occurred in CON.

At posttest, participants in STRUC had significantly increased their functional performance (+ 6%) and their 6-minute walk distance (+ 12%) whereas NEED and CON remained stable. The knee-extension strength significantly decreased (- 6%) from pre- to posttest in participants in CON.

Discussion

STRUC was more effective in improving functional performance and exercise capacity. However, this study provides evidence for the short-term effectiveness of a need-supportive PA program on moderate-intensity PA. Considering cost-effectiveness, the limited number of contact moments in NEED might be more efficient for large scale PA promotion in assisted living facilities. Moreover, need-supportive coaching is hypothesized to enhance autonomous motivation, possibly resulting in more long-term engagement in PA.

Key words

Physical activity, cycle ergometer, intervention, assisted living facilities